

## Environmental Factors that Affect Range Plant Growth

[Llewellyn L. Manske](#) PhD

Range Scientist, NDSU, Dickinson Research Extension Center

Environmental factors affect range plant growth. The three most ecologically important environmental factors affecting rangeland plant growth are light, temperature, and water (precipitation). Plant growth and development are controlled by internal regulators that are modified according to environmental conditions. A research project was conducted to describe the three most important environmental factors in western North Dakota and to identify some of the conditions and variables that limit range plant growth. Rangeland managers should consider these factors during the development of long-term management strategies (Manske 2000).

Light is the most important ecological factor affecting plant growth. Light is necessary for photosynthesis, and changes in day length (photoperiod) regulate the phenological development of rangeland plants. Changes in the day length function as the timer or trigger that activates or stops physiological processes initiating growth and flowering and that starts the process of hardening for resistance to low temperatures in fall and winter. The tilt of the earth's axis in conjunction with the earth's annual revolution around the sun produces the seasons and changes the length of daylight in temperate zones. Dickinson ([Fig. 1](#)) has nearly uniform day and night lengths (12 hours) during only a few days, near the vernal and autumnal equinoxes, 20 March and 22 September, respectively, when the sun's apparent path crosses the equator as the sun travels north or south, respectively. The shortest day length (8 hours, 23 minutes) occurs at winter solstice, 21 December, when the sun's apparent path is farthest south of the equator. The longest day length (15 hours, 52 minutes) occurs at summer solstice, 21 June, when the sun's apparent path is farthest north of the equator. The length of daylight changes during the growing season, increasing from about 13 hours in mid April to nearly 16 hours in mid June, then decreasing to around 11 hours in mid October ([Fig. 1](#)).

Temperature, an approximate measurement of the heat energy available from solar radiation, is a significant factor because both low and high temperatures limit plant growth. Most plant biological activity and growth occur within only a narrow range of temperatures, between 32EF (0EC) and 122EF (50EC). The long-term (108-year) mean annual temperature in the Dickinson, North Dakota, area is 40.7EF (4.8EC) ([Table 1](#)). January is the coldest month, with a mean temperature of 10.9EF (-11.7EC). July and August are the warmest months, with mean temperatures of 68.6EF (20.3EC) and 66.9EF (19.4EC), respectively. Months with mean monthly temperatures below 32.0EF (0.0EC) are too cold for active plant growth. Low temperatures define the growing season for perennial plants, which is generally from mid

April to mid October (6.0 months). Perennial grassland plants are capable of growing for longer than the frost-free period, but to continue active growth, they require temperatures above the level that freezes water in plant tissue and soil. Winter dormancy in perennial plants is not total inactivity but reduced activity.

Water (precipitation) is essential for all plants and is an integral part of living systems. Water is ecologically important because it is a major force in shaping climatic patterns and biochemically important because it is a necessary component in physiological processes. Plant water stress limits growth. Water stress can vary in degree from a small decrease in water potential to the lethal limit of desiccation. The long-term (108-year) annual precipitation for the area of Dickinson, North Dakota, is 16.03 inches (407.09 mm). The growing season precipitation (April to October) is 13.56 inches (344.58 mm) and is 84.54% of the annual precipitation. June has the greatest monthly precipitation, at 3.55 inches (90.22 mm). The seasonal distribution of precipitation ([Table 2](#)) shows the greatest amount of precipitation occurring in the spring (7.30 inches, 45.54%) and the smallest amount occurring in winter (1.54 inches, 9.61%). Total precipitation for November through March averages less than 2.5 inches (15.35%). The precipitation received in May, June, and July accounts for 50.53% of the annual precipitation (8.10 inches).

Of the past 108 years (1892 to 1999), 14 (12.96%) were drought years, receiving 75% or less of the long-term mean precipitation level. Fifteen (13.89%) were wet years, receiving 125% or more of the long-term mean precipitation level. Seventy-nine years (73.15%) received normal annual precipitation amounts, between 75% and 125% of the long-term mean. Of the past 108 growing seasons, 17 (15.74%) were drought growing seasons, 19 (17.59%) were wet growing seasons, and 72 (66.67%) received precipitation at normal levels.

Temperature and precipitation act together to affect the physiological and ecological status of range plants. The balance between rainfall and potential evapotranspiration determines a plant's biological situation. When rainfall is lower than evapotranspiration demand, a water deficiency exists. The ombrothermic graph technique (Emberger et al. 1963), which plots mean monthly temperature and monthly precipitation on the same axis, was used to identify months with water deficiency conditions during 1892-1999 (Manske 2000). The long-term ombrothermic graph for the Dickinson area ([Fig. 2](#)) shows near water deficiency conditions for August, September, and October, a finding indicating that range plants generally may have difficulty growing and accumulating biomass during these 3 months. Favorable water relations occur during May, June, and July, a period during which range plants should be capable of growing and accumulating herbage biomass.

Drought years occurred during 13.0% of the past 108 years, and 15.7% of the growing seasons were drought growing seasons. The 108-year period (1892 to 1999) contained a total of 648 growing season months. Water deficiency conditions occurred during 211.5 of these, a finding indicating that during 32.64% of the growing season months, or for an average of 2.0 months during every 6.0-month growing season, range plants were under water stress and therefore limited in growth and herbage biomass accumulation. Water deficiency occurred in May and June 14.8% and 9.3 % of the time, respectively. Water deficiency conditions occurred in July less than 40% of the time. Water deficiency conditions occurred in August, September, and October more than 50% of the time: 50.0% of the time in August, 50.9 % of the time in September, and 49.1% of the time in October. Water deficiency conditions lasting a month or more cause plants to experience water stress severe enough to reduce herbage production. These levels of water stress are a major factor limiting the quantity and quality of plant growth in western North Dakota and can limit livestock production if not considered during the development and implementation of long-term grazing management strategies.

## Acknowledgment

I am grateful to Amy M. Kraus and Naomi J. Thorson for assistance in preparation of this manuscript. I am grateful to Sheri Schneider for assistance in production of the tables and figures and for word processing this manuscript.

Table 1. Long-term (1892-1999) mean monthly temperature and monthly precipitation at Dickinson, ND.

	<b>B<sub>F</sub></b>	<b>B<sub>C</sub></b>	<b>in.</b>	<b>mm</b>
<b>Jan</b>	10.87	-11.74	0.42	10.63
<b>Feb</b>	15.03	-9.43	0.41	10.30
<b>Mar</b>	25.94	-3.37	0.71	18.06
<b>Apr</b>	41.49	5.27	1.43	36.36
<b>May</b>	52.85	11.58	2.32	58.95
<b>Jun</b>	62.04	16.69	3.55	90.22
<b>Jul</b>	68.55	20.31	2.23	56.54
<b>Aug</b>	66.88	19.38	1.77	45.04
<b>Sep</b>	55.92	13.29	1.32	33.54
<b>Oct</b>	43.85	6.58	0.94	23.93
<b>Nov</b>	28.25	-2.09	0.53	13.54
<b>Dec</b>	16.91	-8.38	0.39	9.99

	<b>MEAN</b>		<b>TOTAL</b>	
	<b>40.72</b>	<b>4.84</b>	<b>16.03</b>	<b>407.09</b>

Table 2. Seasonal percentage of mean annual precipitation distribution (1892-1999).		
<b>Season</b>	<b>in.</b>	<b>%</b>
<b>Winter (Jan, Feb, Mar)</b>	1.54	9.61
<b>Spring (Apr, May, Jun)</b>	7.30	45.5
<b>Summer (Jul, Aug, Sep)</b>	5.32	33.19
<b>Fall (Oct, Nov, Dec)</b>	1.87	11.67
<b>TOTAL</b>	<b>16.03</b>	

Fig. 1. Annual pattern of daylight duration at Dickinson, North Dakota.





Fig. 2. Ombrothermic diagram of long-term (1892-1999) mean monthly temperature and monthly precipitation at Dickinson, North Dakota.



Literature Cited

**Emberger, C., H. Gaussen, M. Kassas, and A. dePhilippis. 1963.** Bioclimatic map of the Mediterranean Zone, explanatory notes. UNESCO-FAO. Paris. 58p.

**Manske, L.L. 2000.** Environmental factors to consider during planning of management for range plants in the Dickinson, North Dakota, region, 1892-1999. NDSU Dickinson Research Extension Center. Range Research Report DREC 00-1018c. Dickinson, ND. 36p.

